

The illogic of Kt/V

Kidney International (2009) **75**, 337; doi:10.1038/ki.2008.584

To the Editor: In reference to the recent articles regarding Kt/V ,^{1,2} a hemodialysis $Kt/V > 1.0$ implies there is complete removal of urea from a volume of body water which exceeds the total volume of water in the body, an impossibility, occurring because it measures dialyzer, not patient, clearance.

Kt/V for peritoneal dialysis is based on the actual amount of urea removed, explaining the discrepancy between the recommended weekly Kt/V in hemodialysis patients, 3.6 (3×1.2)–4.2 (3×1.4), compared to the weekly Kt/V in peritoneal dialysis patients, 1.7–2.0.³ If the urea reduction ratio (URR) represents the actual amount of urea removed per hemodialysis treatment, with three sessions in a week, each with a urea reduction ratio of 0.65–0.70, would remove about 1.95–2.1 times the total body urea content, nearly identical to the recommended weekly Kt/V for peritoneal dialysis.

Kt/V is a flawed concept as used in hemodialysis and results in a fictitious and questionably meaningful number, but does have a useful role in peritoneal dialysis. URR more accurately reflects solute removal.

1. Spalding EM, Chandna SM, Davenport A *et al.* Kt/V underestimates the hemodialysis dose in women and small men. *Kidney Int* 2008; **74**: 348–355.
2. Lowrie EG. Prescribing and monitoring hemodialysis dose. *Kidney Int* 2008; **74**: 262–264.
3. Hemodialysis Adequacy 2006 Work Group. Clinical practice guidelines for hemodialysis adequacy, update 2006. *Am J Kidney Dis* 2006; **48**(Suppl 1): S2–S175.

Paul G. Jenkins¹

¹Milwaukee Kidney Associates, Milwaukee, USA

Correspondence: Paul G. Jenkins, Milwaukee Kidney Associates, 1575 RiverCenter Drive, Suite 104, Milwaukee, Wisconsin 53212, USA.

E-mail: pgjmd@sbcglobal.net

Response to 'The illogic of Kt/V '

Kidney International (2009) **75**, 337; doi:10.1038/ki.2008.586

I agree completely with Dr Jenkins' impression that Kt/V is clinically illogical for reasons explained in the Commentary he cites.¹ In addition, Eloot *et al.* have recently found better urea removal when patients are treated for a longer time (t) despite comparable Kt/V .² Those data illustrate one more reason why K , t , and V should not be combined as a single ratio; we have one more fact indicating the clinical 'illogic of Kt/V '.

I respectfully disagree with his reason, however, because $Kt/V > 1.0$ does not indicate complete urea removal. The fact is revealed by his claim that urea reduction ratio (URR) reflects solute removal better than Kt/V . Urea kinetic equations prove that URR and Kt/V are tightly linked as the following illustrates:

- (1) BUN at end of dialysis (C_t) = predialysis concentration (C_0) $\times e^{-Kt/V}$, so

$$(2) C_t/C_0 = e^{-Kt/V}, \text{ and}$$

$$(3) \text{URR} = 100 \times (C_0 - C_t)/C_0 = 100 \times (1 - C_t/C_0), \text{ so}$$

$$(4) \text{URR} = 100 \times (1 - e^{-Kt/V}).$$

Therefore, URR and Kt/V can be calculated from each other and one cannot reasonably argue in the same breath that URR is appropriate but Kt/V is not.

This example shows that $Kt/V = 0.5 \rightarrow \text{URR} = 39\%$, $Kt/V = 1.0 \rightarrow \text{URR} = 63\%$, and $Kt/V = 1.5 \rightarrow \text{URR} = 78\%$. Note also that $\text{URR} = 99\% \leftarrow Kt/V = 4.6$, $\text{URR} = 99.9\% \leftarrow Kt/V = 6.9$, and $\text{URR} = 99.99\% \leftarrow Kt/V = 9.2$. So, although Kt/V is not appropriate for evaluating clinical outcome,¹ it does reflect that URR and $Kt/V > 1.0$ is perfectly logical.

The confusion probably evolved from misleading statements suggesting Kt/V reflects a 'fractional clearance' of urea. For example, we find,

"To normalize for differences in the size and habitus of patients, a dose of hemodialysis (prescribed or delivered) is best described as the *fractional clearance* of urea as a function of its distribution volume (Kt/V)".³ (Emphasis added)

Such unclear statements likely suggested to many that when, for example, $Kt/V = 48\text{ l}/40\text{ l} = 1.2$ the 'fractional clearance' would be 120%, so those 40 l should be completely 'cleared'. Unfortunately, the concept is incorrect as I hope I have shown.

1. Lowrie EG. Prescribing and monitoring hemodialysis dose. *Kidney Int* 2008; **74**: 262–264.
2. Eloot S, Van Biesen W, Dhondt A *et al.* Impact of hemodialysis duration on the removal of uremic retention solutes. *Kidney Int* 2008; **73**: 765–770.
3. NKF K/DOQI Guidelines 2000, Guideline #2: Method of Measurement of Delivered Dose of Hemodialysis (Evidence) http://www.kidney.org/PROFESSIONALS/kdoqi/guidelines_updates/doqi_uptoc.html.

Edmund G. Lowrie¹

¹Shore Road, Cape Neddick, Maine, USA

Correspondence: Edmund G. Lowrie, 511 Shore Road, Cape Neddick, Maine 3902, USA. E-mail: edlowrie@prodigy.net

Response to 'The illogic of Kt/V '

Kidney International (2009) **75**, 337–338; doi:10.1038/ki.2008.589

Our paper¹ was not intended to praise the Kt/V concept of dialysis adequacy but to provide some constructive criticism. Jenkins raises some theoretical issues that relate to the validity of the whole Kt/V concept in the hemodialysis setting, whereas our paper drew attention to potential problems with the concept of using V , an estimate of total body water, to normalize estimated urea clearance. We provided some support for the use of a normalizing factor, which better reflects metabolic activity.

There are a number of issues. First, whether any measure of urea removal is sufficient to judge dialysis

adequacy; second, if a measure of urea removal were to be used, whether a simple urea reduction ratio is acceptable or whether a measure that incorporates a normalizing factor is necessary; and finally – whether the widespread use of V as this normalizing factor is valid. Jenkins' criticisms relate to the second point whereas our paper addressed the third.

Focusing solely on normalized urea clearance presents a flawed view of dialysis adequacy. There are many other facets. Middle molecule removal is important. Reanalysis of data from the HEMO study has demonstrated that retention of β_2 microglobulin was associated with increased mortality.² In addition to solute clearances, dialysis prescription encompasses other goals, including sodium and water, divalent ion and acid-base homeostasis. Dialysis hypotension is more likely to occur when dialysis session time is shortened, because of increased ultrafiltration rates, and failure to achieve adequate sodium removal with increased interdialytic weight gains.

Although we have criticized the appropriateness of the use of Kt/V in prescribing hemodialysis, we do believe that an assessment of the amount of dialysis delivered is vital, unless very long and/or very frequent dialysis sessions are employed. Normalized urea clearance, although not perfect, is a useful marker of adequacy, though we believe it could be used more astutely. Our paper proposes that the normalizing factor should reflect metabolic activity, as use of V may risk under-dialysis in women and small men. We suggest that this should be borne in mind while we continue to use Kt/V , and that the adequacy targets should be adjusted in these two groups.

1. Spalding EM, Chandna SM, Davenport A *et al.* Kt/V underestimates the hemodialysis dose in women and small men. *Kidney Int* 2008; **74**: 348–355.
2. Cheung AK, Rocco MV, Yan G *et al.* Serum beta-2 microglobulin levels predict mortality in dialysis patients: results of the HEMO study. *J Am Soc Nephrol* 2006; **17**: 546–555.

Elaine M. Spalding¹, Shahid M. Chandna¹, Andrew Davenport² and Ken Farrington¹

¹Renal Unit, Lister Hospital, Stevenage, Herts, UK and ²Renal Unit, Royal Free Hospital, London, UK

Correspondence: Ken Farrington, Renal Unit, Lister Hospital, Coreys Mill Lane, Stevenage, Herts SG1 4AB, UK. E-mail: ken.farrington@nhs.net

Lipid disorders in experimental chronic kidney disease: a role for SREBPs

Kidney International (2009) **75**, 338; doi:10.1038/ki.2008.592

To the Editor: We have read with interest the article by Hai-Lu Zhao *et al.*¹ on fat redistribution and adipocyte transformation in uninephrectomized rats. Among numerous interesting findings, the authors demonstrate increased

ectopic fat deposits in remnant kidneys and other solid organs of chronic kidney disease animals. The authors speculate on the possible mechanisms, taking into account increased lipid production. This hypothesis is strengthened by demonstrating that expression of HMG-CoA reductase, the rate-limiting enzyme in cholesterologenesis pathway, is increased in remnant kidneys.

We would like to confirm these results and highlight the potential role of sterol regulatory element-binding proteins (SREBPs) in the above-mentioned disturbances. SREBPs are nuclear transcription factors that are currently regarded as the major regulators of both cholesterologenesis and lipogenesis.

We have shown that both gene expression and protein abundance of SREBPs are increased in white adipose tissue² and livers³ of chronic kidney disease rats. Moreover, increase in SREBP expression has been demonstrated in kidneys of diabetic mice⁴ and in experimental age-related nephropathy,⁵ where it has been clearly linked to lipid deposition in glomeruli, and consequently to mesangial expansion, glomerulosclerosis, and proteinuria.

These results complement the findings demonstrated by Zhao *et al.* as they bring us closer to elucidating the issue of altered lipid metabolism in the course of chronic kidney disease.³

1. Zhao HL, Sui Y, Guan J *et al.* Fat redistribution and adipocyte transformation in uninephrectomized rats. *Kidney Int* 2008; **74**: 467–477.
2. Karczyska J, Stelmanska E, Nogalska A *et al.* Upregulation of lipogenic enzymes genes expression in white adipose tissue of rats with chronic renal failure is associated with higher level of sterol regulatory element binding protein-1. *Metabolism* 2004; **53**: 1060–1065.
3. Szolkiewicz M, Chmielewski M, Nogalska A *et al.* The potential role of sterol regulatory element binding protein transcription factors in renal injury. *J Ren Nutr* 2007; **17**: 62–65.
4. Sun L, Halaihel N, Zhang W *et al.* Role of sterol regulatory element-binding protein 1 in regulation of renal lipid metabolism and glomerulosclerosis in diabetes mellitus. *J Biol Chem* 2002; **277**: 18919–18927.
5. Jiang T, Liebman SE, Lucia MS *et al.* Role of altered renal lipid metabolism and the sterol regulatory element binding proteins in the pathogenesis of age-related renal disease. *Kidney Int* 2005; **68**: 2608–2620.

Michał Chmielewski¹, Marek Szolkiewicz¹ and Bolesław Rutkowski¹

¹Department of Nephrology, Transplantology and Internal Medicine, Medical University of Gdansk, Gdansk, Poland

Correspondence: Michał Chmielewski, Department of Nephrology, Transplantology and Internal Medicine, Medical University of Gdansk, ul. Debinki 7, 80–211 Gdansk, Poland. E-mail: chmiel@amg.gda.pl

Response to 'Lipid disorders in experimental chronic kidney disease: a role for SREBPs'

Kidney International (2009) **75**, 338–339; doi:10.1038/ki.2008.594

We thank Chmielewski *et al.* for drawing attention to the potential role of sterol regulatory element-binding